

Comprehensive Analysis of Antarctic Ozone Hole Dynamics Using Day- Night IASI Infrared Observations

Guido Masiello^a, Tiziano Maestri^b, Carmine Serio^a, Giuliano Liuzzi^a, Angela Cersosimo^a, Michele Martinazzo^b, Rocco Giosa^a, Lorenzo Cassini^{a,c}, Federico Donat^{a,b}, Fabrizio Masin^b, Pamela Pasquariello^a, Giorgia Proietti Pelliccia^b, Sara Venafra^d, Claude Camy-Peyret^e

- a. DiIng, Università degli studi della Basilicata, Potenza, Italy
- b. DIFA, Università di Bologna, Italy
- c. DICEA, University of Rome, Italy
- d. Italian Space Agency, Rome, Italy
- e. Institut Pierre-Simon Laplace, Paris, France

Anknowledgements

- This work is supported by Italian Space Agency in the framework of the following project
 - FORUM-Scienza (contract n-2018-6-Q.0, consortium led by CNR-INO Florence),
 - Fit-FORUM (contract n. 2023-23-HH.0, CUP F33C23000240005, consortium led by DIFA, University of Bologna),
 - MC-FORUM (contract n. 2023-23-HH.0, CUP F93C2300046000, consortium led by IBE-CNR)



Agenzia
Spaziale
Italiana



Foreword

- 2021 and 2023 have been two years with the most spatially extensive, and deep ozone hole.
- Formation of **Polar stratospheric clouds (PSCs)** is the fundamental catalytic mechanism that accelerates ozone destruction
- PSC formation involves **HNO₃** and H₂O initially in the gas phase, which condenses into the solid phase (giving rise to crystals of HNO₃-3H₂O or NAT) at **T<195 K**.
- The phenomenon is continuously monitored by satellite instruments, (Ozone Monitoring Instrument, OMI, TROPOspheric Monitoring Instrument, TROPOMI)
- They need daylight
- They have no sensitivity to the thermodynamic conditions of UT/LS region, and they don't sense nitric acid and water in the gas phase.

Outline




DATA & METHOD



RESULTS



CONCLUSIONS





The “Ingredients” of the Ozone Depletion

	TROPOMI	OMI	IASI	MLS-AURA
Ozone	X	X	X	
HNO ₃			X	X
Temperature			X	
Water Vapour			X	

We collect data for the 9th of July, Sept and Oct 2021 and 2023 $-90^{\circ}<Lat<-60^{\circ}$

Microwave Limb Sounder (MLS) on NASA's EOS AURA Satellite



- Level 3 observations of the HNO₃ profiles for MLS.
- <https://disc.gsfc.nasa.gov/>
- Sept. 9, 2021, and 2023.
- Level 3 data available on a 4°×5° longitude/latitude grid
- pressure range: 215–1.5 hPa.
- HNO₃ Profiles → Columnar Amount

$$\bar{X}_{HNO_3} = \frac{1}{p_u - p_l} \int_{p_l}^{p_u} q(p) dp$$

OMI and TROPOMI

TROPospheric
Monitoring Instrument
(TROPOMI) Level 2
data (flag $q < 0.75$)

Ozone Measurements
Instruments (OMI)
Level 3 gridded data

φ -IASI-F2N: forward/inverse package from the far to near infrared




φ -IASI-F2N is a forward/inverse for all-sky (clear and cloudy conditions) calculations of forward (L1) and inverse (L2) products

The system includes

- σ -IASI-F2N
- δ -IASI-F2N

To sum up

- φ stands for φυσικά physical
- σ stands for wavenumber and for s - Spectrum
- δ stands for increment





Serio et al. 2024 JQSRT

δ -IASI/F2N State Vector


Atmosphere (profiles of size $N_L=60$)		Surface
• Temperature (K)	• CH ₄ (ppv)	• Ts (scalar, K)
• H ₂ O (g/kg)	• SO ₂ (ppv)	• Emissivity spectrum, $\varepsilon(\sigma)$
• HDO (ppv)	• NH ₃ (ppv)	Cloud parameters (profiles of size, $N_L=60$)
• CO ₂ (ppv)	• HNO ₃ (ppv)	• LWC (kg/kg)
• O ₃ (ppv)	• CF ₄ (ppv)	• IWC (kg/kg)
• N ₂ O (ppv)	• OCS (ppv)	• r_e (μm)
• CO (ppv)		• D_e (μm)
		• Cloud fraction, cf (scalar)

$$\mathbf{v} = \left(\begin{array}{c} \varepsilon(\sigma), cf, T_s, \mathbf{T}, \mathbf{Q}, \mathbf{O}, \mathbf{HDO}, \mathbf{q}_{CO_2}, \\ \mathbf{q}_{CH_4}, \mathbf{q}_{N_2O}, \mathbf{q}_{CO}, \mathbf{q}_{SO_2}, \mathbf{q}_{NHO_3}, \mathbf{q}_{HNO_3}, \mathbf{q}_{OCS}, \mathbf{q}_{CF_4}, \\ \mathbf{q}_{LWC}, \mathbf{q}_{IWC}, \mathbf{q}_{r_e}, \mathbf{q}_{D_e}, w \end{array} \right)$$

1 day IASI B & C Antarctica overpasses




Retrieval of Polar Stratospheric Cloud




Orange – IASI
Measurement
Black – Computed
with ECMWF profile
Blue – Computed
with IASI retrieved
profile

Retrieval of Polar Stratospheric Cloud



$P=43 \text{ hPa}$ ($\sim 20 \text{ km}$)
 $T=182 \text{ K}$
 $\text{IWC}=4.5 \cdot 10^{-7} \text{ kg/kg}$
 $D_e=4 \mu \text{m}$


Retrieval of Ozone, 09/09/2021



- a) O₃ map from ECMWF (background),
- b) O₃ map retrieved by IASI data (un-gridded level 2 product)

➤ IASI see a deeper and wider Ozone hole with respect ECMWF


Retrieval of Ozone, 09/09/2021




- a) O₃ map from ECMWF (background),
 - b) O₃ map retrieved by IASI data (un-gridded level 2 product)
 - c) OMI level 3 gridded and smoothed product.
 - d) TROPOMI level 2 gridded product, (QF>0.75)
- IASI see a deeper and wider Ozone hole with respect ECMWF
 - OMI and TROPOMI don't cover the pole (Night)

2021 Total Ozone zonal mean

- OMI and TROPOMI don't cover the pole (Night)
- IASI see a deeper and wider Ozone hole with respect ECMWF.
- Overall, our IASI level 2 data compare better with TROPOMI than OMI.
- OMI tends to be more coherent with the ECMWF analysis, could be just the results that OMI total ozone is of the level 3 type.





P_{\min} , T_{\min} & HNO_3






- In the inner continent, the temperature is well below **195 K** (b)
- Pressure at Temperature inversion ranges 30-60 hPa (a)
- Low amount of HNO_3 < 500 pptv in the inner continent (c)
- PSCs formation may lead to the removal of nitric acid from the gas phase.

Zonal Mean of T_{\min} & HNO_3




HNO₃ (IASI v/s MLS)





Zonal average
of on 9 Sept.
2021;
left y-axis IASI;
right y-axis
MLS_aura

Red – IASI
Blue – FIT
Black -FG



Magenta-noise
Blue – FIT
Black - FG



Spectral
Residuals:
Cloudy
over Land

Key Points

- **ϕ -IASI-F2N:** we developed a system integrating a direct and inverse models. The scheme simultaneously retrieve cloud optical properties, the atmosphere's thermodynamic, and composition state for the clear-cloudy sky
- **Infrared Instrument Importance:** Infrared instruments like IASI are crucial for improving the night/day coverage and spatial resolution of trace gases, enhancing our understanding of ozone depletion in the polar atmosphere.
- **Ozone Hole and Denitrification:** The onset of the ozone hole in September 2021 and 2023 (not shown here) was accompanied by significant denitrification of the polar atmosphere.
- **Instrument Agreement:** Both IASI (infrared) and MLS_aura (microwave) instruments observed this denitrification, despite their different viewing geometries and spectral ranges.
- **NAT/PSC Clouds:** NAT PSC clouds catalyzed ozone destruction, contributing to the overall ozone hole.